arXiv:1304.5900v1 [math.AG] 22 Apr 2013

A short note on cubic fourfolds containing a plane and K3 surfaces of Picard rank two

Federica Galluzzi *

Abstract

We present some families of cubic hypersurfaces in $\mathbb{P}^5(\mathbb{C})$ containing a plane whose associated quadric bundle does not have a rational section.

Mathematics Subject Classification (2010): 14E08,14C30,14J28,14F22.

1 Introduction

Let X be a smooth cubic hypersurface in $\mathbb{P}^5(\mathbb{C})$. Investigating the rationality of X is a classical problem in algebraic geometry. The general X is conjectured to be not rational but not a single example of non rational cubic fourfold is known. Cubic fourfolds containing a quartic scroll or a quintic del Pezzo surface are rational (see [F], [Mo]). Idem for those fourfolds containing a plane and a Veronese surface (see [Tr]). Beauville and Donagi showed in [BD] that also pfaffian cubic fourfolds are rational.
The cubic fourfolds containing a plane form a divisor $\mathcal{C}_8$ in the moduli space $\mathcal{C}$ of all cubic fourfolds. The general fourfold containing a plane is also expected to be non rational. Nevertheless, Hassett showed in [H1] that there exists a countable infinite collection of divisors in C_8 which parameterize rational cubic fourfolds. The fourfolds containing a plane are birational to the total space of a quadric surface bundle by projecting from the plane: Hassett's examples are rational since the associated quadric bundle has a rational section.
Recently, Auel at al. (see [ABBV]) have described a class of rational cubic fourfolds whose associated quadric bundle do not have a rational section. The general member is rational since it is in an irreducible component of $\mathcal{C}_8 \cap \mathcal{C}_{14}$ where $\mathcal{C}_{14}$ denotes the divisor of $\mathcal{C}$ containing pfaffian cubic fourfolds.
Using results on the Hodge structure of cubic fourfolds and $K3$ surfaces, we present a family of cubic fourfolds containing a plane whose associated quadric bundle does not have a rational section. We don't know if these fourfolds are rational. The rational examples in [ABBV] are members of our family.

The paper is organized as follows. In Sections 2 and 3 we recall some basic notions on lattices and $K3$ surfaces. We focus on $K3$ surfaces of Picard rank

---

[*] The author is supported by the framework PRIN 2010/11 "Geometria delle Varietà Algebriche", cofinanced by MIUR.

two recalling the fundamental work of Nikulin in [N]. Then in 3.1 we present the $K3$ surfaces of Picard rank two which are double covers of the plane ramified over a sextic curve. In 3.1.1 we construct a family $S_{(d,c)}$ of double planes with Picard rank two. In Section 4 we recall how these surfaces are related to cubic $4-$folds containing a plane. A such cubic X is birational to a quadric bundle $Y \xrightarrow{\pi} \mathbb{P}^2$ which, in the general case, ramifies over a smooth sextic curve C. The Hodge structure of X is strictly related to the Hodge structure of the $K3$ surface S obtained as a double cover of the plane ramified over C and parameterizing the rulings of the quadrics in the fibration $Y \xrightarrow{\pi} \mathbb{P}^2$ (see [V, §1]). We use the following fact: the lattice $A(X)$ of $2-$cycles modulo numerical equivalence on X has rank three and even discriminant if S has Picard rank two and even Néron-Severi discriminant (see [V, Proposition 2] and [ABBV, Proposition 1,Corollary 2]). In case of $\mathrm{rk}(A(X)) = 3$ we prove that the quadric bundle $Y \xrightarrow{\pi} \mathbb{P}^2$ does not have a rational section if and only if the discriminant of $A(X)$ is even (see 4.0.7).

In 4.1 we recover the cubic hypersurfaces associated to the double planes $S_{(d,c)}$ using the additional datum of an odd theta characteristic on the discriminant sextic (see [B2, V]).

Finally, in Theorem 4.1.2 we prove that the quadric bundles associated to those hypersurfaces with d even do not have a rational section. The rational examples in [ABBV] correspond to fourfolds associated to $S_{(2,-1)}$.

Acknowledgments. The author would like to thank Edoardo Sernesi and Bert van Geemen for helpful remarks and useful hints.

2 Lattices

A *lattice* is a free $\mathbb{Z}$-module L of finite rank with a $\mathbb{Z}$-valued symmetric bilinear form $b_L(x, y)$. A lattice is called *even* if the quadratic form associated to the bilinear form has only even values, *odd* otherwise. The *discriminant* $d(L)$ of a lattice is the determinant of the matrix of its bilinear form. A lattice is called *non-degenerate* if the discriminant is non-zero and *unimodular* if the discriminant is ± 1. If the lattice L is non-degenerate, the pair (s_+, s_-), where $s_\pm$ denotes the multiplicity of the eigenvalue ± 1 for the quadratic form associated to $L \otimes \mathbb{R}$, is called *signature* of L. Finally, we call $s_+ + s_-$ the *rank* of L and L is said *indefinite* if the associate quadratic form has both positive and negative values. Given a lattice L, the lattice $L(m)$ is the $\mathbb{Z}$-module L with bilinear form $b_{L(m)}(x, y) = m b_L(x, y)$. An *isometry* of lattices is an isomorphism preserving the bilinear form. Given a sublattice $L' \subset L$, the embedding is *primitive* if $\frac{L}{L'}$ is free. Recall the formula

$$[L : L']^2 = \frac{d(L')}{d(L)}, \tag{1}$$

where $L' \subset L$ is a sublattice with $\mathrm{rk}(L') = \mathrm{rk}(L)$.

2.1 Examples.

i) The lattice $\langle n \rangle$ is a free $\mathbb{Z}$-module of rank one , $\mathbb{Z}\langle e \rangle$, with bilinear form $b(e,e) = n$.

ii) The *hyperbolic lattice* is the even, unimodular, indefinite lattice with $\mathbb{Z}$-module $\mathbb{Z}\langle e_1, e_2 \rangle$ and bilinear associated form given by the matrix $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. We write
$$U = \left\{ \mathbb{Z}^2, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right\}.$$

iii) The lattice E_8 has $\mathbb{Z}^8$ as $\mathbb{Z}$-module and the matrix of the bilinear form is the Cartan matrix of the root system of E_8. It is an even, unimodular and positive definite lattice.

3 K3 surfaces of rank two

A $K3$ surface is a smooth projective surface S with trivial canonical class and $H^1(S, \mathcal{O}_S) = 0$. It is well known that $H^2(S, \mathbb{Z})$ is an even, unimodular and positive definite lattice, with respect to the intersection form on S. It has rank 22, signature $(3, 19)$ and it is isomorphic to
$$\Lambda := U^{\oplus 3} \oplus E_8(-1)^{\oplus 2}.$$
The lattice Λ will be called the $K3$ *lattice*. The Hodge numbers are $(1, 20, 1)$, (see [BPV, VIII]). Denote with
$$N(S) \cong H^2(S, \mathbb{Z}) \cap H^{1,1}(S)$$
the *Néron-Severi lattice* of S, it is a primitive sublattice. If S is a $K3$ surface and
$$c_1 : Pic(S) \longrightarrow H^2(S, \mathbb{Z})$$
is the first Chern class, one has $Pic(S) \cong c_1(Pic(S)) = NS(S)$.
The orthogonal complement $T(S)$ of $N(S)$ in $H^2(S, \mathbb{Z})$ is the *transcendental lattice* of S.
The *rank* of S, $\rho(S)$, is the rank of $N(S)$. The Hodge Index Theorem implies that $N(S)$ has signature $(1, \rho(S) - 1)$ and that $T(S)$ has signature $(2, 20 - \rho(S))$.
Let $l \in NS(S)$ be a class with $l^2 > 0$. The *primitive cohomology* $H^2(S, \mathbb{Z})_0$ is the orthogonal complement of the lattice $< l >$.
Main tools for the study of K3 surfaces are Torelli theorem (see [LP] and [PSS]) and the Surjectivity of the Period Map (see [T]). The *period* of S is given by $[\omega_S] = \mathbb{P}(H^{2,0}(S))$ in the period domain
$$\Omega = \{x \in \mathbb{P}(\Lambda \otimes \mathbb{C}) | x \cdot x = 0, \, x \cdot \bar{x} > 0\} \subset \mathbb{P}(\Lambda \otimes \mathbb{C}).$$
By Torelli Theorem and the surjectivity of the period map, the period determines the $K3$ surface: given $\alpha \in \Omega$ there exists a $K3$ surface S_α (unique up to isomorphism) with period α such that $H^2(S_\alpha, \mathbb{Z})$ is isometric to Λ.

Nikulin in [N] made a deep study of lattice theory and of integral quadratic forms and use them to study $K3$ surfaces. We recall the following which is crucial for our purposes

Theorem 3.0.1. *[N, Theorem 1.14.4] [M, Corollary 2.9] If* $\rho(S) \leq 10$, *then every even lattice* M *of signature* $(1, \rho - 1)$ *occurs as the Néron-Severi group of some* $K3$ *surface and the primitive embedding* $M \hookrightarrow \Lambda$ *is unique.*

Corollary 3.0.2. *All even lattices of rank* 2 *and signature* $(1,1)$ *occur as the Néron-Severi lattice* $N(S)$ *of some K3-surface* S *of rank two and the primitive embedding* $N(S) \hookrightarrow \Lambda$ *is unique. A generic such lattice has the form*

$$M = \left\{ \mathbb{Z}^2, \begin{pmatrix} 2a & b \\ b & 2c \end{pmatrix} \right\}.$$

3.1 $K3$ surfaces double planes of rank two

A double covering of the projective plane $\varphi : S \longrightarrow \mathbb{P}^2$ branched along a smooth sextic C is a $K3$ surface: $\varphi_*(\mathcal{O}_S) \cong \mathcal{O}_{\mathbb{P}^2} \oplus \mathcal{O}(3)$, so $H^1(S, \mathcal{O}_S) = 0$ and $\omega_S \cong \varphi^*(\omega_{\mathbb{P}^2} \otimes \mathcal{O}(3)) \cong \mathcal{O}_S$. The surface S is smooth since C is. Moreover, two surfaces of this type are birationally equivalent if and only if they have the same ramification divisor. A class $l \in NS(S)$ with $l^2 > 0$ is the pull-back of the class of a line in $\mathbb{P}^2$ and here $l^2 = 2$. If S has rank two the Néron-Severi lattice has the form

$$L_{(b,c)} = \left\{ \mathbb{Z}^2, \begin{pmatrix} 2 & b \\ b & 2c \end{pmatrix} \right\}.$$

The $K3$ surface S in this case is called a *double plane.* For general references on double planes, see [En] and [S].

3.1.1 Examples.

i) Consider S a $K3$ surface double plane ramified over a smooth sextic with Néron-Severi lattice of the form

$$L_{(1,-1)} = \left\{ \mathbb{Z}^2, \begin{pmatrix} 2 & 1 \\ 1 & -2 \end{pmatrix} \right\}.$$

This can be realized taking a double copver of the plane ramified over a sextic curve having a tritangent line l. The pull-back of l to S is a divisor splitting into two irreducible components l_1, l_2 . The corresponding divisor classes are linearly independent. Both curves are isomorphic to l and $l_1^2 = l_2^2 = -2$.

ii) Analogously, if the Néron-Severi lattice has the form

$$L_{(2,-1)} = \left\{ \mathbb{Z}^2, \begin{pmatrix} 2 & 2 \\ 2 & -2 \end{pmatrix} \right\}$$

the corresponding double plane S can be realized with a ramification sextic C which is tangent to a conic D in 6 points with multiplicity two. As before, $\varphi^*(D) = D_1 + D_2,$ with D_1, D_2 isomorphic to D and $D_1^2 = D_2^2 = -2$.

The previous examples can be generalized as follows.

Lemma 3.1.1. *The lattice*

$$L_{(d,c)} = \left\{ \mathbb{Z}^2, \begin{pmatrix} 2 & d \\ d & 2c \end{pmatrix} \right\}$$

is the Néron-Severi lattice of a $K3$ *surface* $S_{(d,c)}$ *whose ramification curve has an everywhere tangent curve of degree* d.

Proof. By Theorem 3.0.1 the lattice $L_{(d,c)}$ can be embedded in Λ as a primitive sublattice. Consider the $K3$ surface $S_{(d,c)} = S_\alpha$ defined by $\alpha \in \Omega$ with $\alpha^\perp = L_{(d,c)}$ and generic with this property; hence $L_{(d,c)} = NS(S_{(d,c)})$. The surface $S_{(d,c)}$ is a double plane. Take a divisor $D \in Pic(S_{(d,c)})$ such that $D \notin |nl|$ for every n. One has $\varphi^*(\varphi(D)) = D_1 + D_2$ and $\varphi(D)$ is everywhere tangent to the ramification sextic. $\square$

4 Cubic $4-$folds containing a plane

Let X be a smooth cubic hypersurface in $\mathbb{P}^5(\mathbb{C})$. Consider the cohomology group $H^4(X, \mathbb{Z})$ and denote with

$$A(X) = H^4(X, \mathbb{Z}) \cap H^{2,2}(X)$$

the lattice of the middle integral cohomology Hodge classes. Those classes are algebraic since X verifies the integral Hodge conjecture (see [Mu] and [Zu]). The *transcendental lattice* $T(X)$ is the orthogonal complement of $A(X)$ (with respect to the intersection form on X).

Suppose now that X contains a plane P. Consider the projection from P onto a plane in $\mathbb{P}^5$ disjoint from P. Blowing up X along P one obtains a quadric bundle $\pi : Y \longrightarrow \mathbb{P}^2$ whose discriminant curve is a plane sextic C. If X does not contain a second plane intersecting P, the curve C is smooth and this means that the quadrics of the bundle have rank ≥ 3 (see [V, Lemme 2]).

Denote with Q the class of such a quadric. One has $P + Q = H^2$, where H is the hyperplane class associated to the embedding $X \hookrightarrow \mathbb{P}^5(\mathbb{C})$. The hypersurface X is said to be *very general* if $A(X) = < H^2, P >$ $(= < H^2, Q >)$. Denote $L := < H^2, P >^\perp$.

X is rational iff the quadric bundle $\pi : Y \longrightarrow \mathbb{P}^2$ is rational and a sufficient condition for the rationality of Y is the existence of a rational section. This last condition may be translated in a condition on the parity of the intersection of some $2-$cycles on X. More precisely, for a $2-$cycle T in X consider the intersection index

$$\delta(T) = T \cdot Q.$$

Note that $\delta(P) = -2$ and $\delta(H^2) = 2$ So, if X is very general the index δ takes only even values. There is the following

Theorem 4.0.2. *([H2, Theorem 3.1.]) If X is a cubic fourfold containing a plane, then $\pi : Y \longrightarrow \mathbb{P}^2$ has a rational section if and only if there exists a cycle T in $A(X)$ with $\delta(T)$ odd.*

Using this Theorem it is easy to give (lattice-theoretic) hints to construct cubic fourfolds with $rk(A(X)) > 2$ whose associated quadric bundles do not have a rational section.

Proposition 4.0.3. *Let X be a smooth cubic fourfold containing a plane P and let $A(X)$ be of rank three. If the the discriminant of $A(X)$ is even the corresponding quadric bundle $\pi : Y \longrightarrow \mathbb{P}^2$ does not have a rational section.*

Proof. The quadric bundle $\pi : Y \longrightarrow \mathbb{P}^2$ has a rational section if and only if there exists a cycle $T \in A(X)$ such that $\delta(T) = c$ is odd (by Theorem 4.0.2). Suppose that such T exists and consider the lattice $< H^2, Q, T > \subset A(X)$. The intersection matrix is given by

$$\begin{array}{c|ccc} & H^2 & Q & T \\ \hline H^2 & 3 & 2 & a \\ Q & 2 & 4 & c \\ T & a & c & b \end{array} \tag{2}$$

for some $a, b, c \in \mathbb{Z}$. Moreover c is odd if and only if the discriminant $-4a^2+8b+4ca-3c^2$ is. In this case the discriminat of $A(X)$ must be odd, contradiction. (See also [ABBV, Proposition 3]). $\square$

Our aim now is to build some geometric examples. To do this, we need to better understand the links between Hodge theory and the geometry on a cubic $4-$fold containing a plane. Here we follow Voisin [V, §1].

Let $\varphi : S \longrightarrow \mathbb{P}^2$ be the double plane branched over C, discriminant sextic of the quadric bundle $Y \longrightarrow \mathbb{P}^2$. The surface S parameterizes the rulings of the quadrics of the fibration. Let F be the Fano variety of lines in X, the subvariety of the Grassmannian $Gr(1,5)$ parameterizing lines contained in X. The divisor $D \subset F$ consisting of lines meeting P is identified with

$$D = \{(l, s) \in F \times S \ : \ l \text{ is in the ruling of the quadric parameterized by } \varphi(s)\}.$$

giving a $\mathbb{P}^1-$bundle

$$f : D \longrightarrow S. \tag{3}$$

The incidence graph restricted to D

$$\begin{tikzcd} D \times X \supset Z_D \arrow[r, "p"] \arrow[d, "q"] & D \\ X & \end{tikzcd}$$

defines the Abel-Jacobi map:

$$\alpha_D = p_* q^* : H^4(X, \mathbb{Q}) \longrightarrow H^2(D, \mathbb{Q})$$

which induces an isomorphism of Hodge structures, see [V, Proposition 1]. There are the following

Proposition 4.0.4. *([V, Proposition 2],[ABBV, Proposition 1]) Let X be a smooth cubic fourfold containing a plane. Then $\alpha_D(L) \subset f^*(H^2(S,\mathbb{Z})_0(-1))$ is a polarized Hodge substructure of index* 2.

Proposition 4.0.5. *[ABBV, Corollary 2] Let X be a smooth cubic fourfold containing a plane. Then $\alpha_D(T(X)) \subset f^*T(S)(-1)$ is a sublattice of index ϵ dividing* 2*. In particular,* $\operatorname{rk} A(X) = \operatorname{rk}(NS(S)) + 1$ *and* $d(A(X)) = 2^{2(\epsilon-1)} d(NS(S))$.

This implies the following

Proposition 4.0.6. *Let X be a cubic fourfold containing a plane. If the bundle $\pi : Y \longrightarrow \mathbb{P}^2$ does not have a rational section then $d(A(X))$ is even.*

Proof. The $\mathbb{P}^1$-bundle $f : D \longrightarrow S$ in (3) produces an element α of order two in the Brauer group $Br(S)$ of S. Recall that, if S is a $K3$ surface, then

$$Br(S) \cong T(S)^* \otimes \mathbb{Q}/\mathbb{Z} \cong \operatorname{Hom}(T(S), \mathbb{Q}/\mathbb{Z})$$

(see for example [vG, §2.1.]). An element α of order 2 defines a surjective homomorphism

$$\alpha : T(S) \longrightarrow \mathbb{Z}/2\mathbb{Z}$$

and thus defines a sublattice T_α of index 2 in $T(S)$ (see [vG, §2.4.]). The quadric bundle associated to X does not have a rational section if and only if α is not trivial in $Br(S)$ (see [Ku, Proposition 4.7.]). If α is not trivial, then $\alpha_D(T(X)) \subset f^*T(S)(-1)$ is a sublattice of index 2 and thus $d(A(X))$ is even by Proposition 4.0.5. □

Corollary 4.0.7. *Let X be a cubic fourfold containing a plane with* $\operatorname{rk}(A(X)) = 3$*. The bundle $\pi : Y \longrightarrow \mathbb{P}^2$ does not have a rational section if and only if $d(A(X))$ is even.*

Proof. It follows by Proposition 4.0.3 and Proposition 4.0.6

□

4.1 Theta-characteristics on the ramification curve C

A theta-characteristic on a smooth curve C is a line bundle κ such that $\kappa^{\otimes 2} = K_C$. We write $h^0(\kappa) := \dim H^0(C, \kappa)$.

Denote with Q_x a quadric of the bundle $Y \longrightarrow \mathbb{P}^2$. The map $x \mapsto Q_x \cap P$ gives a net of conics whose discriminant curve is a plane cubic C_1. The curve C_1 cuts a divisor $2D$ on the sextic C and thus it determines an effective theta-characteristic on C (see [V, Lemme 7]). Conversely, the cubic hypersurface X is determined by the curve C plus an odd theta-characteristic (see [V, Proposition 4]). The same result is implied by the following

Proposition 4.1.1. *([B2, Proposition 4.2.]) Let C be a smooth plane curve of degree d, defined by an equation $F = 0$ and κ an odd theta-characteristic on C with $h^0(\kappa) = 1$. Thus, κ admits a minimal resolution*

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}^2}(-2)^{d-3} \oplus \mathcal{O}_{\mathbb{P}^2}(-3) \xrightarrow{M} \mathcal{O}_{\mathbb{P}^2}(-1)^{d-3} \oplus \mathcal{O}_{\mathbb{P}^2} \longrightarrow \kappa \longrightarrow 0$$

with a symmetric matrix $M \in M_{(d-2)\times(d-2)}(\mathbb{C}[X_0, X_1, X_2])$ satisfying $\det M = F$, *and of the form*

$$M = \begin{pmatrix} L_{1,1} & \dots & L_{1,d-3} & Q_1 \\ \vdots & & \vdots & \vdots \\ L_{1,d-3} & \dots & L_{d-3,d-3} & Q_{d-3} \\ Q_1 & \dots & Q_{d-3} & H \end{pmatrix} \tag{4}$$

where the forms $L_{i,j}$, Q_i, H are linear, quadratic and cubic respectively. Conversely, the cokernel of a symmetric matrix M as above is an odd theta-characteristic κ on C with $h^0(\kappa) = 1$.

We can now prove our main result.

Theorem 4.1.2. *Consider the couple $(S_{(d,c)}, \kappa)$ where $S_{(d,c)}$ is a double plane defined as in Lemma 3.1.1 and κ is a theta characteristic on the ramification curve C with $h^0(\kappa) = 1$. If d is even, $(S_{(d,c)}, \kappa)$ determines a cubic fourfold containing a plane whose associated quadric bundle does not have a rational section.*

Proof. Let C be the ramification curve of $S := S_{(d,c)}$ and take a theta characteristic κ on C with $h^0(\kappa) = 1$. Proposition 4.1.1 says that the curve C has a determinantal representation $F = \det M = 0$ with

$$M = \begin{pmatrix} L_{1,1} & L_{1,2} & L_{1,3} & Q_1 \\ L_{1,2} & L_{2,2} & L_{2,3} & Q_2 \\ L_{1,3} & L_{2,3} & L_{3,3} & Q_3 \\ Q_1 & Q_2 & Q_3 & H \end{pmatrix}.$$

The geometric interpretation is the following. Choose projective coordinates $[Z_1, Z_2, Z_3, X_0, X_1, X_2]$ in $\mathbb{P}^5(\mathbb{C})$ and define the cubic fourfold $X = X(S, \kappa)$ as the zero set

$$\sum_{i,j=1}^{3} Z_iZ_jL_{i,j}(X_0,X_1,X_2)+\sum_{i=1}^{3} 2Z_iQ_i(X_0,X_1,X_2)+H(X_0,X_1,X_2)=0.$$

The cubic X is smooth and it contains the plane $P := X_0 = X_1 = X_2 = 0$. The curve C is the discriminant of the quadric bundle obtained by projecting the hypersurface X from P.

The $K3$ surface S has rank two and d is even, so the discriminant of $NS(S)$ is even. This means that $A(X)$ has rank three and even discriminant by Proposition 4.0.5. Thus Proposition 4.0.3 implies that the corresponding quadric bundle $\pi : Y \longrightarrow \mathbb{P}^2$ does not have a rational section. □

Remark 4.1.3. Auel et al. in [ABBV] (see Proposition 4 and Theorem 11) construct an explicit example of a pfaffian (then rational) cubic fourfold associated to a $K3$ surface of type $S_{(2,-1)}$.

References

[ABBV] A. Auel, M. Bernadara, M. Bolognesi, A. Várilly-Alvarado , *Cubic fourfolds containing a plane and a quintic del Pezzo surface*, arXiv:math.AG/1205.0237v3.

[BPV] W. Barth, C. Peters, A. Van de Ven, Compact complex surfaces, Ergebnisse der Mathematik und ihrer Grenzgebiete, Springer-Verlag,2004.

[B1] A. Beauville, *Variétés de Prym et jacobiennes intermédiaires*, Ann. Sci. Ec. Norm. Sup. **10**, (1977), 309-391.

[B2] A. Beauville, *Determinantal hypersurfaces*, Michigan Math. J. **48**, no.1 (2000), 39-64.

[BD] A. Beauville, R. Donagi, *La variété des droites d'une hypersurface cubique de dimension* 4, C. R. Acad. Sci. Paris Ser. I Math., **301** (1985), 703-706.

[En] F. Enriques, *Sui piani doppi di genere uno*, Memorie della Società Italiana delle Scienze, s. III, t. X (1896), 201-222.

[F] G. Fano, *Sulle forme cubiche dello spazio a cinque dimensioni contenenti rigate del quarto ordine*, Comment. Math. Helv. **15** (1943), 71-80.

[H1] B. Hassett, *Some rational cubic fourfolds*, J. Algebraic Geom. **8** no. 1 (1999), 103-114.

[H2] B. Hassett, *Special cubic fourfolds*, Compositio Math. **120** no.1 (2000), 1-23.

[Ku] A. Kuznetsov, *Derived categories of cubic fourfolds*, in "Cohomological and Geometric Approaches to Rationality Problems. New Perspectives". Progress in Mathematics **282** F. Bogomolov; Y. Tschinkel, (Eds.) 2010.

[LP] E. Looijenga, C. Peters, *A Torelli theorem for K3 surfaces* , Comp. Math. **42** (1981), 145-186.

[Mo] U. Morin , *Sulla razionalità dell'ipersuperficie cubica generale dello spazio lineare a cinque dimensioni*, Rend. Sem. Mat. Univ. di Padova, **11** (1940) 108-112.

[M] D.R. Morrison *On $K3$ surfaces with large Picard number*, Invent. Math. **75** (1984) 105-121.

[Mu] J.P. Murre, *On the Hodge conjecture for unirational fourfolds*, Indag. Math. **39**, no. 3, (1977), 230-232.

[N] V. Nikulin *Integral symmetric bilinear forms and some of their applications*, Math. USSR Izv. **14** (1980) 103-167.

[PSS] I: Piatechki-Shapiro, I.R. Shafarevich , *A Torelli theorem for algebraic surfaces of type K3*, Math. USSR Izvestija **5** (1971), 547-588.

[S] E. Sernesi *Introduzione ai piani doppi*, in Seminario di Geometria 1977/78, Centro di Analisi Globale, Firenze (1979) 1-78.

[T] A.N. Todorov, *Applications of the Kähler-Einstein- Calabi-Yau metric to moduli of K3 surfaces* , Invent. Math. **61** (1980), 251-265.

[Tr] S. L. Tregub, *Three constructions of rationality of a cubic fourfold*, Moscow Univ. Math. Bull. **39** no. 3, (1984), 8-16.

[vG] B. van Geemen, *Some remarks on Brauer groups of K3 surfaces*, Adv. Math. **197**, no. 1, (2005) 222-247.

[V] C. Voisin , *Théorème de Torelli pour les cubiques de P^5. (French) [A Torelli theorem for cubics in $\mathbf{P}^5$]*, Invent. Math. **86**, no. 3, (1986), 577-601.

[Zu] S. Zucker, *The Hodge conjecture for cubic fourfolds*, Compositio Math., **34**, no. 2, (1977), 199-209.

Federica Galluzzi
Dipartimento di Matematica
Università di Torino
Via Carlo Alberto n.10 ,Torino
10123, ITALIA
e-mail: `federica.galluzzi@unito.it`